



Korlym[®]
mifepristone
300 mg Tablets

YOUR TREATMENT. YOUR JOURNEY.

A guide for patients taking **Korlym** that includes helpful tips and tools to support you throughout treatment.

INDICATION

Korlym is a prescription medicine used to treat high blood sugar (hyperglycemia) caused by high cortisol levels in the blood (hypercortisolism) in adults with endogenous Cushing's syndrome who have type 2 diabetes mellitus or glucose intolerance and have failed surgery or cannot have surgery.

Korlym is not for people who have type 2 diabetes mellitus not caused by Cushing's syndrome.

It is not known if **Korlym** is safe and effective in children.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about Korlym?

Korlym can cause serious side effects, including:

- **Loss of a pregnancy.** Women who can become pregnant must:
 - have a negative pregnancy test before starting **Korlym** or before restarting **Korlym** if you stop taking it for more than 14 days
 - use a non-hormonal form of birth control while taking **Korlym** and for one month after stopping **Korlym**. Talk to your doctor about how to prevent pregnancy. Tell your doctor right away if you think you may be pregnant.

Please see additional Important Safety Information throughout, and [Important Facts](#) about Korlym.

Quick contacts

My Healthcare Professional(s)

Name: _____

Specialty: _____

Phone: _____

Name: _____

Specialty: _____

Phone: _____

My Corcept Cares Patient Advocate

Name: _____

Phone: _____

Phone: **1-855-295-9969**

Email: **patientadvocate@corcept.com**

Your **Patient Advocate** can only reach out to you with your consent. Please call the **Patient Advocate** number or Specialty Pharmacy number below to give your consent for **Patient Advocate** support.

My Specialty Pharmacy

1-855-4KORLYM (1-855-456-7596)

A Specialty Pharmacist is on call 24 hours a day, 7 days a week to answer your questions about taking **Korlym** and other medications.

To learn more about the support program, see [*Help at every step along the way.*](#)
For more information, visit [Korlym.com](https://www.korlym.com).

Your journey begins here

Treatment tips	5
Prepare for your journey	7
Cortisol and your body	9
What is Korlym?	13
What can I expect with Korlym?	15
Side effects of Korlym	17
What is the best dose for me?	19
Check-ins along the way	21
Help at every step along the way	23
Support through Specialty Pharmacies	25
Your Patient Access Coordinator	26
Preparing for your next appointment	27
Important Safety Information	29

Please see additional Important Safety Information throughout, and [Important Facts](#) about Korlym.

Korlym®
mifepristone
300 mg Tablets



Now that you have been diagnosed with Cushing syndrome (hypercortisolism), you can take steps toward improving some of your symptoms. Remember everyone's journey is different.

This guide will help you understand what to expect along the way and where to find support when you need it.

IMPORTANT SAFETY INFORMATION (CONT.)

Do not take Korlym if you:

- are pregnant
- are taking medicines that contain simvastatin, lovastatin, cyclosporine, dihydroergotamine, ergotamine, fentanyl, pimozide, quinidine, sirolimus, or tacrolimus
- must take corticosteroid medicines for other serious medical problems
- have experienced unexplained vaginal bleeding, changes in the cells lining your uterus (endometrial hyperplasia), or cancer of the lining of your uterus (endometrial cancer)
- are allergic to mifepristone or any of the ingredients in **Korlym**.

Please see additional Important Safety Information throughout, and Important Facts about Korlym.

Korlym®
mifepristone
300 mg Tablets

Treatment tips



Be patient
as your body adjusts
to treatment.



Set goals
with the help of
your healthcare
professional team.



**Tell your healthcare
professional about any
side effects.**

You may need to adjust one
or more of your medications.
Certain side effects can be expected
early in treatment and may decrease
as you find the best dose.



Ask for help
when you need it.
Your **Patient Advocate**
can provide support
and information.*

*Although **Patient Advocates** provide personalized support, they cannot offer medical advice. Talk regularly with your healthcare professional about your treatment and any side effects you may experience.

IMPORTANT SAFETY INFORMATION (CONT.)

What Are the Possible Side Effects of Korlym?

Serious side effects of Korlym may also include:

- **reduced effects of adrenal hormones (adrenal insufficiency).** Tell your doctor right away if you have unusual tiredness or weakness, nausea, fatigue, low blood pressure (hypotension), low blood sugar (hypoglycemia)

Please see additional Important Safety Information throughout, and Important Facts about Korlym.

Korlym®
mifepristone
300 mg Tablets

Treatment tips



Plan a regular time to take your Korlym dose every day.
Korlym should be taken once daily with a meal.



Korlym tablets should be swallowed whole.
Do not split, crush, or chew the tablets.



Get involved.
Ask questions about your treatment.



Use the checklists.
They can help you prepare for appointments.
(See [Appointment checklist](#))



Use the progress tracker.
Discuss it with your healthcare professional at your appointments.
(See [Progress tracker](#))



Look forward,
but don't forget to look back now and then to see how far you've come.
The progress tracker can help with that, too!

IMPORTANT SAFETY INFORMATION (CONT.)

What Are the Possible Side Effects of Korlym?

Serious side effects of Korlym may also include (cont.):

- **low blood potassium (hypokalemia).** Your doctor should check the level of potassium in your blood before you start taking **Korlym** and while you take it. Tell your doctor if you have muscle weakness, aches or cramps, abnormal or irregular heartbeats (palpitations)

Please see additional Important Safety Information throughout, and [Important Facts](#) about Korlym.

Korlym®
mifepristone
300 mg Tablets

Prepare for your journey

Before treatment begins

You will have testing done to evaluate your blood sugar, blood pressure, and potassium levels.

Work with your healthcare professional to develop personalized treatment goals and understand what you can expect during treatment, including anticipated side effects. Discuss any monitoring you will need to do, as well as medication adjustments or discontinuation.

» My treatment goals

Write down the treatment goals you set with your healthcare professional.

Potassium levels should be checked before starting treatment, 1 to 2 weeks after the first dose, 1 to 2 weeks after a dose increase, and periodically as treatment continues. Blood sugar and blood pressure should be monitored regularly throughout treatment.

IMPORTANT SAFETY INFORMATION (CONT.)

What Are the Possible Side Effects of Korlym?

Serious side effects of Korlym may also include (cont.):

- **bleeding from the vagina.** Korlym may cause the lining of your uterus to become thick and may cause your uterus to bleed. Tell your doctor right away about any bleeding from your vagina that is not normal for you
- **disruption of menstrual cycle**

Please see additional Important Safety Information throughout, and [Important Facts](#) about Korlym.

Korlym®
mifepristone
300 mg Tablets

Prepare for your journey

Achieving your treatment goals may take time. Here is what to expect throughout your treatment with **Korlym**.

Early in treatment

In the first 6 weeks, you may see changes in your:

- » Blood sugar levels, including a reduction in dose of diabetes medications (such as insulin)
- » Mood*
- » Ability to think clearly*

Some of the most common side effects occur during the first 2 weeks of treatment as your body adjusts to less cortisol activity.

Tell your healthcare professional about any side effects you experience and reach out to your **Patient Advocate** when you need support.

To learn more, see [Side effects of Korlym](#).

Later in treatment

After 2 months, you may continue to see improvements in your:

- » Blood sugar levels, including a reduction in dose of diabetes medications (such as insulin)

You may also see improvements in your*:

- » Weight
- » Skin (such as less acne)
- » Body hair (hair growth patterns)
- » Stretch marks
- » Other signs of endogenous hypercortisolism (such as fat deposits or rounded face)

Work with your healthcare professional to assess your progress and manage any side effects you are experiencing. Your healthcare professional will adjust your dose as needed.

*In the clinical study, some patients receiving **Korlym** reported changes in weight, waist measurement, appearance (such as the clearing of acne), mood, symptoms of depression, ability to think clearly, and muscle strength. It is uncertain whether **Korlym** caused these changes. At the start of the trial, patients had an average weight of 220 pounds and an average waistline of 47 inches.

Please see additional information about the trial on [Page 15](#).

IMPORTANT SAFETY INFORMATION (CONT.)

What Are the Possible Side Effects of Korlym?

Serious side effects of Korlym may also include (cont.):

- problems with the electrical system of your heart (QT interval prolongation)
- worsening of symptoms of other medical problems that are treated with corticosteroids when you take corticosteroids and Korlym at the same time.

Please see additional Important Safety Information throughout, and [Important Facts](#) about Korlym.

Korlym[®]
mifepristone
300 mg Tablets

Cortisol and your body

Cushing syndrome (hypercortisolism) is a complex disease that can be difficult to understand. Getting to know the basics of cortisol and its effects on your body is an important part of your treatment journey.

What is cortisol?

Cortisol is a natural hormone produced by your body. It helps to regulate blood sugar, blood pressure, and other important bodily functions.

IMPORTANT SAFETY INFORMATION (CONT.)

What Are the Possible Side Effects of Korlym? (cont.)

The most common side effects of Korlym include nausea, fatigue, headache, low potassium in your blood, pain in your arms and legs (arthralgia), vomiting, swelling of your arms and legs (peripheral edema), high blood pressure, dizziness, decreased appetite, and thickening of the lining of the uterus (endometrial hypertrophy).

These are not all the possible side effects of **Korlym**. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please see additional Important Safety Information throughout, and [Important Facts](#) about Korlym.

Korlym[®]
mifepristone
300 mg Tablets

Cortisol and your body



What is hypercortisolism?

Hypercortisolism occurs when your body makes too much cortisol on its own. Endogenous hypercortisolism occurs when a growth on the pituitary gland, adrenal glands, or elsewhere in the body causes the production of too much cortisol. This increase in cortisol activity can result in high blood sugar, weight gain, infection, hypertension, and cardiovascular disease.

Cushing syndrome is caused by excess cortisol—also called hypercortisolism.

IMPORTANT SAFETY INFORMATION (CONT.)

Before taking Korlym, tell your doctor if you are pregnant or plan to become pregnant, have low potassium in your blood (hypokalemia), have or have had a bleeding problem or are taking medicines to thin your blood, have or have had heart problems, have had an organ transplant, have been taking medicines called corticosteroids (cortisone, dexamethasone, methylprednisolone, prednisolone, prednisone), are breastfeeding or plan to breastfeed.

You should not drink grapefruit juice while you take **Korlym**.

Tell your healthcare provider about all the medicines you take, including prescription and nonprescription medicines, vitamins, and herbal supplements. Using **Korlym** with certain other medicines may affect each other or may cause serious side effects.

Please see additional Important Safety Information throughout, and Important Facts about Korlym.

Korlym[®]
mifepristone
300 mg Tablets

Cortisol and your body

What are the signs and symptoms of hypercortisolism?

Hypercortisolism is a challenging disease that disrupts your body and mind—even when you're doing all you can to lead a healthy lifestyle.

The signs and symptoms of hypercortisolism vary from one person to the next, so the problems you are facing could be very different from others with hypercortisolism.

IMPORTANT SAFETY INFORMATION (CONT.)

What is the most important information I should know about Korlym?

Korlym can cause serious side effects, including:

- **Loss of a pregnancy.** Women who can become pregnant must:
 - have a negative pregnancy test before starting **Korlym** or before restarting **Korlym** if you stop taking it for more than 14 days
 - use a non-hormonal form of birth control while taking **Korlym** and for one month after stopping **Korlym**. Talk to your doctor about how to prevent pregnancy. Tell your doctor right away if you think you may be pregnant.

Please see additional Important Safety Information throughout, and Important Facts about Korlym.

Korlym[®]
mifepristone
300 mg Tablets

Cortisol and your body

Hypercortisolism can look different for everyone. Here are some of the ways it can impact your body and mental well-being.



Health

- » Obesity or weight gain
- » High blood sugar or diabetes
- » High blood pressure
- » Irregular periods
- » Muscle weakness
- » Weak bones
- » Hardening of the arteries
- » Slow wound healing
- » Frequent infections
- » Fluid retention, swelling of feet/legs
- » Kidney stones



Mood/Energy

- » Mood changes: depression, anxiety, irritability
- » Poor short-term memory
- » Poor concentration
- » Low sex drive
- » Low energy, fatigue/tiredness
- » Trouble sleeping



Appearance

- » Rounded face
- » Unwanted hair growth, hair loss
- » Red or flushed face
- » Belly fat
- » Extra fat around neck
- » Skin: red or purple stripes or streaks, stretch marks, easy bruising, acne

IMPORTANT SAFETY INFORMATION (CONT.)


Do not take Korlym if you:

- are pregnant
- are taking medicines that contain simvastatin, lovastatin, cyclosporine, dihydroergotamine, ergotamine, fentanyl, pimozide, quinidine, sirolimus, or tacrolimus
- must take corticosteroid medicines for other serious medical problems
- have experienced unexplained vaginal bleeding, changes in the cells lining your uterus (endometrial hyperplasia), or cancer of the lining of your uterus (endometrial cancer)
- are allergic to mifepristone or any of the ingredients in **Korlym**.

Please see additional Important Safety Information throughout, and Important Facts about Korlym.

Korlym
mifepristone
300 mg Tablets

What is **Korlym**?



Your healthcare professional has prescribed **Korlym** for you. **Korlym** is a medicine that can improve certain signs and symptoms of endogenous Cushing syndrome.

So, how does it work?

IMPORTANT SAFETY INFORMATION (CONT.)

What Are the Possible Side Effects of Korlym?

Serious side effects of Korlym may also include:

- **reduced effects of adrenal hormones (adrenal insufficiency).** Tell your doctor right away if you have unusual tiredness or weakness, nausea, fatigue, low blood pressure (hypotension), low blood sugar (hypoglycemia)

Please see additional Important Safety Information throughout, and Important Facts about Korlym.

Korlym[®]
mifepristone
300 mg Tablets

What is **Korlym**?



Korlym is a prescription medicine used to treat high blood sugar (hyperglycemia) caused by high cortisol levels in the blood (hypercortisolism) in adults with endogenous Cushing syndrome who have type 2 diabetes mellitus or glucose intolerance and have failed surgery or cannot have surgery.

Korlym is not for people who have type 2 diabetes mellitus not caused by Cushing syndrome.

It is not known if **Korlym** is safe and effective in children.

IMPORTANT SAFETY INFORMATION (CONT.)

What Are the Possible Side Effects of Korlym?

Serious side effects of Korlym may also include (cont.):

- **low blood potassium (hypokalemia)**. Your doctor should check the level of potassium in your blood before you start taking **Korlym** and while you take it. Tell your doctor if you have muscle weakness, aches or cramps, abnormal or irregular heartbeats (palpitations)

Please see additional Important Safety Information throughout, and Important Facts about Korlym.

Korlym[®]
mifepristone
300 mg Tablets

What can I expect with **Korlym**?

At first, your treatment with **Korlym** may be challenging. Your body has been affected by high cortisol activity for a long time. Over time, your body will adjust to decreased cortisol activity. Some improvements will happen quickly and others will take time.

Clinical changes with **Korlym**

Korlym was studied in a 24-week clinical trial of 50 adults with endogenous Cushing syndrome who could not have surgery or for whom surgery had failed. All patients received **Korlym**. **Korlym's** effect on blood sugar was tested in 29 patients with type 2 diabetes or impaired blood sugar control.*

In patients with type 2 diabetes or impaired blood sugar control

- » **60% (15/25) of patients saw at least a 25% improvement in their blood sugar levels**
 - » Some patients saw improvement in blood sugar within 6 weeks after starting **Korlym**
- » **47% (7/15) of patients taking antidiabetic medicine had their medication reduced**

*Patients were also assessed based on 8 broad categories: glucose control, lipids, blood pressure, body composition, clinical appearance, strength, psychiatric/cognitive symptoms, and quality of life.

IMPORTANT SAFETY INFORMATION (CONT.)

What Are the Possible Side Effects of Korlym?

Serious side effects of Korlym may also include (cont.):

- **bleeding from the vagina.** **Korlym** may cause the lining of your uterus to become thick and may cause your uterus to bleed. Tell your doctor right away about any bleeding from your vagina that is not normal for you
- **disruption of menstrual cycle**

Please see additional Important Safety Information throughout, and [Important Facts](#) about **Korlym**.

Korlym[®]
mifepristone
300 mg Tablets

What can I expect with Korlym?



Clinical changes with **Korlym** (continued)

Across the entire study group

- » **Patients taking Korlym lost an average of 5.7% of their body weight after 6 months***
 - » On average, men (14 patients) lost 3.3 inches and women (32 patients) lost 2.7 inches from their waistline
- » **Patients reported an overall decrease in depression symptoms***

Cortisol withdrawal

It can take time for your body to adjust to less cortisol activity after starting Korlym. During this period, you might experience something known as "cortisol withdrawal."

This may include experiencing flu-like symptoms. Some symptoms of cortisol withdrawal may include feeling very tired, aches or weakness in your muscles or joints, nausea, and abdominal pain. These are not all of the possible side effects of **Korlym**. These symptoms may improve as you reach the best dose of **Korlym** for your body.

*In the clinical study, some patients receiving **Korlym** reported changes in weight, waist measurement, appearance (such as the clearing of acne), mood, symptoms of depression, ability to think clearly, and muscle strength. It is uncertain whether **Korlym** caused these changes. At the start of the trial, patients had an average weight of 220 pounds and an average waistline of 47 inches.

IMPORTANT SAFETY INFORMATION (CONT.)

What Are the Possible Side Effects of Korlym?

Serious side effects of Korlym may also include (cont.):

- problems with the electrical system of your heart (QT interval prolongation)
- worsening of symptoms of other medical problems that are treated with corticosteroids when you take corticosteroids and Korlym at the same time.

Please see additional Important Safety Information throughout, and Important Facts about Korlym.

Korlym[®]
mifepristone
300 mg Tablets

Side effects of **Korlym**

It's common to have some side effects as your body gets used to less cortisol activity. Tell your healthcare professional about any signs or symptoms you experience.

The most common side effects of Korlym include:

- Nausea
- Fatigue
- Headache
- Low potassium in your blood
- Pain in your arms and legs (arthralgia)
- Vomiting
- Swelling of your arms and legs (peripheral edema)
- High blood pressure
- Dizziness
- Decreased appetite
- Thickening of the lining of the uterus (endometrial hypertrophy)

Cortisol withdrawal

If your body has been exposed to high cortisol activity over time, it may have gotten used to it. **It can take time to adjust to less cortisol activity after starting Korlym. During this period, you might experience "cortisol withdrawal."** Some symptoms include fatigue (feeling very tired), nausea, aches or weakness in your muscles or joints, and abdominal pain. These are not all of the possible side effects of **Korlym**. If you experience these symptoms, talk to your healthcare professional.

Changes in blood sugar or blood pressure*

When you regularly monitor your blood sugar and blood pressure at home, record the results. Let your healthcare professional know of any change in your blood sugar or blood pressure levels. It's important to know if either of these levels is higher or lower than normal. Your healthcare professional may need to adjust your medications. (See [Check-ins along the way](#) and [Progress tracker](#))

***Korlym** is not indicated to treat high blood pressure.

IMPORTANT SAFETY INFORMATION (CONT.)

What Are the Possible Side Effects of Korlym? (cont.)

The most common side effects of Korlym include nausea, fatigue, headache, low potassium in your blood, pain in your arms and legs (arthralgia), vomiting, swelling of your arms and legs (peripheral edema), high blood pressure, dizziness, decreased appetite, and thickening of the lining of the uterus (endometrial hypertrophy).

These are not all the possible side effects of **Korlym**. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA.

Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please see additional Important Safety Information throughout, and [Important Facts](#) about Korlym.

Korlym
mifepristone
300 mg Tablets

Side effects of **Korlym**

It's possible to experience serious side effects during treatment with **Korlym**.

Unusual vaginal bleeding

If you have a uterus and are taking **Korlym**, you may have unusual vaginal bleeding, such as spotting, missed periods, irregular periods, or other menstrual changes. If you notice signs of unusual bleeding, tell your healthcare professional right away. It is also recommended that you notify your gynecologist that you are taking **Korlym**.

Adrenal insufficiency

If you experience unusual tiredness, weakness, nausea, or fatigue accompanied by low blood pressure and/or low blood sugar, contact your healthcare professional right away.

Low potassium levels (hypokalemia)

Muscle weakness, aches, cramps, abnormal or irregular heartbeat (palpitations), constipation, and swelling of the hands or feet (edema) could be a sign of low potassium levels. Tell your healthcare professional if you have any of these symptoms while taking **Korlym**. Low potassium while you are taking **Korlym** should be treated by your healthcare professional.

(See [Check-ins along the way](#) for more information about potassium levels.)

These are not all of the possible side effects of Korlym. Tell your healthcare professional if you have a side effect that bothers you or does not go away.

IMPORTANT SAFETY INFORMATION (CONT.)

Before taking Korlym, tell your doctor if you are pregnant or plan to become pregnant, have low potassium in your blood (hypokalemia), have or have had a bleeding problem or are taking medicines to thin your blood, have or have had heart problems, have had an organ transplant, have been taking medicines called corticosteroids (cortisone, dexamethasone, methylprednisolone, prednisolone, prednisone), are breastfeeding or plan to breastfeed.

You should not drink grapefruit juice while you take **Korlym**.

Tell your healthcare provider about all the medicines you take, including prescription and nonprescription medicines, vitamins, and herbal supplements. Using **Korlym** with certain other medicines may affect each other or may cause serious side effects.

Please see additional Important Safety Information throughout, and [Important Facts](#) about Korlym.

Korlym
mifepristone
300 mg Tablets

What is the best dose for me?

Your healthcare professional will work with you to find the dose that will help you reach your treatment goals. **Korlym** dosing is a process, and it takes time. It's important to understand that not all patients taking **Korlym** need the same dose (number of tablets each day).

In the clinical study of **Korlym**, patients first experienced improvements at different doses. All of the patients started with 1 tablet (300 mg) a day, but most needed a higher dose before they saw results.



Please see additional information about the trial on [Page 15](#).

IMPORTANT SAFETY INFORMATION (CONT.)

What is the most important information I should know about Korlym?

Korlym can cause serious side effects, including:

- **Loss of a pregnancy.** Women who can become pregnant must:
 - have a negative pregnancy test before starting **Korlym** or before restarting **Korlym** if you stop taking it for more than 14 days
 - use a non-hormonal form of birth control while taking **Korlym** and for one month after stopping **Korlym**. Talk to your doctor about how to prevent pregnancy. Tell your doctor right away if you think you may be pregnant.

Please see additional Important Safety Information throughout, and [Important Facts](#) about Korlym.

Korlym
mifepristone
300 mg Tablets

What is the best dose for me?



Korlym should be taken once a day, with a meal, and swallowed whole. **Korlym** should not be split, chewed, or crushed.

Treatment starts with one 300-mg tablet daily.

300
MG/DAY



1 tablet

600
MG/DAY



2 tablets

900
MG/DAY



3 tablets

1200
MG/DAY



4 tablets
Maximum dose

Over time, your healthcare professional may slowly increase your dose 1 tablet at a time, to up to 4 tablets or 1200 mg daily. This will depend on how your body responds to treatment and how your symptoms improve. Your dose will likely not change more often than every 2-4 weeks.

In people with reduced kidney function or those with mild to moderate liver problems, the maximum dose of **Korlym** should be limited to 600 mg/day. Talk to your healthcare professional if you have a history of kidney or liver problems.

IMPORTANT SAFETY INFORMATION (CONT.)

Do not take Korlym if you:

- are pregnant
- are taking medicines that contain simvastatin, lovastatin, cyclosporine, dihydroergotamine, ergotamine, fentanyl, pimozide, quinidine, sirolimus, or tacrolimus
- must take corticosteroid medicines for other serious medical problems
- have experienced unexplained vaginal bleeding, changes in the cells lining your uterus (endometrial hyperplasia), or cancer of the lining of your uterus (endometrial cancer)
- are allergic to mifepristone or any of the ingredients in **Korlym**.

Please see additional Important Safety Information throughout, and Important Facts about Korlym.

Korlym
mifepristone
300 mg Tablets

Check-ins along the way

During your treatment, it's very important that you monitor your blood sugar and blood pressure. Your healthcare professional will also monitor your potassium levels.

Monitoring these levels can help your healthcare professional see how your treatment is working and know when to adjust your **Korlym** dose or doses of other medications you take. Record your results from home testing and lab visits on the [Progress tracker](#). Bring your entries with you to appointments to discuss with your healthcare professional.

Three important checks

✓ 1. Blood sugar

Check your blood sugar regularly. Tell your healthcare professional about any changes.

Why? Changes in your blood sugar are an early and ongoing way to see how your treatment with **Korlym** is working. **Korlym** can help reduce blood sugar due to excess cortisol, so medications you take for diabetes may need to be reduced or adjusted. Your healthcare professional will adjust your prescriptions as needed.

Important: Inform your healthcare professional right away if you have low blood sugar.

✓ 2. Potassium levels

The level of potassium in your blood should be checked:

- Before starting treatment with **Korlym**
- 1 to 2 weeks after your first dose
- 1 to 2 weeks after a dose increase
- Periodically throughout treatment

Why? Because low potassium, called hypokalemia, can happen at any time during treatment with **Korlym**. If you experience muscle weakness, aches, or cramps, abnormal or irregular heartbeat, constipation, or swelling of your hands or feet (edema), it could be due to low potassium (See [Side effects of Korlym](#) to learn more). Work with your healthcare professional to make sure your potassium levels are checked regularly.

IMPORTANT SAFETY INFORMATION (CONT.)

What Are the Possible Side Effects of Korlym?

Serious side effects of Korlym may also include:

- **reduced effects of adrenal hormones (adrenal insufficiency).** Tell your doctor right away if you have unusual tiredness or weakness, nausea, fatigue, low blood pressure (hypotension), low blood sugar (hypoglycemia)

Please see additional Important Safety Information throughout, and [Important Facts](#) about Korlym.

Korlym®
mifepristone
300 mg Tablets

Check-ins along the way

✓ 3. Blood pressure*

Check your blood pressure regularly. Tell your healthcare professional about any changes.

Why? Because if you take blood pressure medication, the dose may need to be adjusted up or down. Your healthcare professional will adjust your prescription as needed.

Important: If you feel dizzy or have other signs of low blood pressure, check your blood pressure and tell your healthcare professional about your signs and symptoms.

***Korlym** is not indicated to treat high blood pressure.

†In the clinical study, some patients receiving **Korlym** reported changes in weight, waist measurement, appearance (such as the clearing of acne), mood, symptoms of depression, ability to think clearly, and muscle strength. It is uncertain whether **Korlym** caused these changes. At the start of the trial, patients had an average weight of 220 pounds and an average waistline of 47 inches.

Other signs of progress†

You may also experience improvements such as:

- » Weight loss and changes in waist measurement
- » Changes in how you look, such as acne clearing up
- » Better mood, fewer symptoms of depression
- » Clearer thinking
- » Improved muscle strength

IMPORTANT SAFETY INFORMATION (CONT.)

What Are the Possible Side Effects of Korlym?

Serious side effects of Korlym may also include (cont.):

- **low blood potassium (hypokalemia).** Your doctor should check the level of potassium in your blood before you start taking **Korlym** and while you take it. Tell your doctor if you have muscle weakness, aches or cramps, abnormal or irregular heartbeats (palpitations)

Please see additional Important Safety Information throughout, and Important Facts about Korlym.

Korlym®
mifepristone
300 mg Tablets

Help at every step along the way

Patient Advocates

Personalized, one-on-one support is available to you during your treatment through a dedicated **Concept Cares Patient Advocate**. Once you have signed up to connect, a **Patient Advocate** will reach out to offer support.



Haven't signed up yet?
Visit [Korlym.com](https://www.korlym.com) or
call 1-855-295-9969
to connect with a
Patient Advocate.

IMPORTANT SAFETY INFORMATION (CONT.)

What Are the Possible Side Effects of Korlym?

Serious side effects of Korlym may also include (cont.):

- **bleeding from the vagina.** Korlym may cause the lining of your uterus to become thick and may cause your uterus to bleed. Tell your doctor right away about any bleeding from your vagina that is not normal for you
- **disruption of menstrual cycle**

Please see additional Important Safety Information throughout, and [Important Facts](#) about Korlym.

Korlym[®]
mifepristone
300 mg Tablets

Help at every step along the way

Useful resources are available to help you succeed in your treatment journey.

How can a Corcept Cares Patient Advocate help me?

Your Patient Advocate is here to



Listen to you when you need someone to talk to about your treatment



Answer your questions about endogenous hypercortisolism, how **Korlym** works, and what to expect during treatment



Remind you to tell your healthcare professional about important things, such as other medications you are taking



Help you understand why it's important to monitor your potassium, blood pressure,* and blood sugar levels



Check in with you to see how you are doing on your current dose



Provide emotional support throughout your treatment with **Korlym**

***Korlym** is not indicated to treat high blood pressure.

*Although **Patient Advocates** provide personalized support, they cannot offer medical advice. Talk regularly with your healthcare professional about your treatment and any side effects you may experience.*



Reach out to a **Corcept Cares Patient Advocate**
1-855-295-9969

IMPORTANT SAFETY INFORMATION (CONT.)

What Are the Possible Side Effects of Korlym?

Serious side effects of Korlym may also include (cont.):

- problems with the electrical system of your heart (QT interval prolongation)
- worsening of symptoms of other medical problems that are treated with corticosteroids when you take corticosteroids and Korlym at the same time.

Please see additional Important Safety Information throughout, and Important Facts about Korlym.

Korlym
mifepristone
300 mg Tablets

Support through Specialty Pharmacies

Korlym is not available at your local retail pharmacy

It is only commercially available through Specialty Pharmacy partners, who will send **Korlym** directly to your home.

Important: **Korlym** cannot be sent to you without your approval. Your Patient Access Coordinator will contact you before every shipment. Be sure to respond!



IMPORTANT SAFETY INFORMATION (CONT.)

What Are the Possible Side Effects of Korlym? (cont.)

The most common side effects of Korlym include nausea, fatigue, headache, low potassium in your blood, pain in your arms and legs (arthralgia), vomiting, swelling of your arms and legs (peripheral edema), high blood pressure, dizziness, decreased appetite, and thickening of the lining of the uterus (endometrial hypertrophy).

These are not all the possible side effects of **Korlym**. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please see additional Important Safety Information throughout, and Important Facts about Korlym.

Korlym
mifepristone
300 mg Tablets

Your Patient Access Coordinator

Support at every step of your journey



Before your first shipment, your Patient Access Coordinator connects with your insurance company.



Then, your Patient Access Coordinator contacts you to schedule your first prescription delivery.



As you progress, your Patient Access Coordinator reaches out to you monthly to schedule refill deliveries.



Reach out to your Specialty Pharmacist if you plan to have any medical procedures, medication changes, or insurance changes.

Your Specialty Pharmacy team is available between 8 AM and 8 PM ET, Monday through Friday.

1-855-4KORLYM (1-855-456-7596)

IMPORTANT SAFETY INFORMATION (CONT.)

Before taking Korlym, tell your doctor if you are pregnant or plan to become pregnant, have low potassium in your blood (hypokalemia), have or have had a bleeding problem or are taking medicines to thin your blood, have or have had heart problems, have had an organ transplant, have been taking medicines called corticosteroids (cortisone, dexamethasone, methylprednisolone, prednisolone, prednisone), are breastfeeding or plan to breastfeed.

You should not drink grapefruit juice while you take **Korlym**.

Tell your healthcare provider about all the medicines you take, including prescription and nonprescription medicines, vitamins, and herbal supplements. Using **Korlym** with certain other medicines may affect each other or may cause serious side effects.

Please see additional Important Safety Information throughout, and Important Facts about Korlym.

Korlym®
mifepristone
300 mg Tablets

Preparing for your next appointment



It's important to keep track of how you are feeling when taking **Korlym**.

Download the [Progress tracker](#) and share your entries with your healthcare professional.

Use the checklist on the next page to prepare for appointments with your healthcare professional. Write down any questions, notes, or reminders in the space provided. This will help them better understand how your body is adjusting to **Korlym**.

Appointment checklist

Before my appointment

- ☐ Review my [Progress tracker](#) and other notes and bring them with me to my appointment
 - ☐ Write down questions I want to ask my healthcare professional
-

During my appointment

- ☐ Discuss
 - Any side effects or new symptoms
 - Ways to manage side effects
 - Any positive changes I have seen
 - Any upcoming medical procedures or surgeries
 - ☐ Ask if my dose should be adjusted
 - ☐ Discuss my other medications and any needed adjustments
 - ☐ Schedule any needed lab tests
 - ☐ Schedule my next visit
-

After my appointment

- ☐ *If there has been a change in any of my medications:*
Call **1-855-456-7596** and talk to a Specialty Pharmacist
 - ☐ Mark my calendar for my next appointment and for any lab tests
 - ☐ Continue to track my symptoms
-

Questions and reminders

INDICATION

Korlym is a prescription medicine used to treat high blood sugar (hyperglycemia) caused by high cortisol levels in the blood (hypercortisolism) in adults with endogenous Cushing's syndrome who have type 2 diabetes mellitus or glucose intolerance and have failed surgery or cannot have surgery.

Korlym is not for people who have type 2 diabetes mellitus not caused by Cushing's syndrome.

It is not known if **Korlym** is safe and effective in children.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about Korlym?

Korlym can cause serious side effects, including:

- **Loss of a pregnancy.** Women who can become pregnant must:
 - have a negative pregnancy test before starting **Korlym** or before restarting **Korlym** if you stop taking it for more than 14 days
 - use a non-hormonal form of birth control while taking **Korlym** and for one month after stopping **Korlym**. Talk to your doctor about how to prevent pregnancy. Tell your doctor right away if you think you may be pregnant.

Do not take Korlym if you:

- are pregnant
- are taking medicines that contain simvastatin, lovastatin, cyclosporine, dihydroergotamine, ergotamine, fentanyl, pimozone, quinidine, sirolimus, or tacrolimus
- must take corticosteroid medicines for other serious medical problems
- have experienced unexplained vaginal bleeding, changes in the cells lining your uterus (endometrial hyperplasia), or cancer of the lining of your uterus (endometrial cancer)
- are allergic to mifepristone or any of the ingredients in **Korlym**.

What Are the Possible Side Effects of Korlym?

Serious side effects of Korlym may also include:

- **reduced effects of adrenal hormones (adrenal insufficiency).** Tell your doctor right away if you have unusual tiredness or weakness, nausea, fatigue, low blood pressure (hypotension), low blood sugar (hypoglycemia)
- **low blood potassium (hypokalemia).** Your doctor should check the level of potassium in your blood before you start taking **Korlym** and while you take it.

Tell your doctor if you have muscle weakness, aches or cramps, abnormal or irregular heartbeats (palpitations)

- **bleeding from the vagina.** **Korlym** may cause the lining of your uterus to become thick and may cause your uterus to bleed. Tell your doctor right away about any bleeding from your vagina that is not normal for you
- **disruption of menstrual cycle**
- **problems with the electrical system of your heart (QT interval prolongation)**
- **worsening of symptoms of other medical problems that are treated with corticosteroids when you take corticosteroids and Korlym at the same time.**

The most common side effects of Korlym include nausea, fatigue, headache, low potassium in your blood, pain in your arms and legs (arthralgia), vomiting, swelling of your arms and legs (peripheral edema), high blood pressure, dizziness, decreased appetite, and thickening of the lining of the uterus (endometrial hypertrophy).

These are not all the possible side effects of **Korlym**. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Before taking Korlym, tell your doctor if you are pregnant or plan to become pregnant, have low potassium in your blood (hypokalemia), have or have had a bleeding problem or are taking medicines to thin your blood, have or have had heart problems, have had an organ transplant, have been taking medicines called corticosteroids (cortisone, dexamethasone, methylprednisolone, prednisolone, prednisone), are breastfeeding or plan to breastfeed.

You should not drink grapefruit juice while you take **Korlym**.

Tell your healthcare provider about all the medicines you take, including prescription and nonprescription medicines, vitamins, and herbal supplements. Using **Korlym** with certain other medicines may affect each other or may cause serious side effects.

Please see [Important Facts](#) about **Korlym** with important warning.

Korlym®
mifepristone
300 mg Tablets



Get 1:1 support throughout your treatment

Corcept Cares Patient Advocates are available to provide personalized support throughout treatment. They can offer educational information and resources on:

- Hypercortisolism
- How **Korlym** works
- Why monitoring is important
- What to expect during treatment

*Although **Patient Advocates** provide personalized support, they cannot offer medical advice. Talk regularly with your healthcare professional about your treatment and any side effects you may experience.*



Connect with a Patient Advocate by scanning the QR code, visiting Korlym.com, or calling 1-855-295-9969.

Please see Important Safety Information throughout, and [Important Facts](#) about Korlym, including an important warning on loss of a pregnancy.