



Talking about Korlym with your healthcare professional team

Conversation guide for patients taking
Korlym® (mifepristone) 300 mg Tablets

Korlym®
mifepristone
300 mg Tablets

INDICATION

Korlym is a prescription medicine used to treat high blood sugar (hyperglycemia) caused by high cortisol levels in the blood (hypercortisolism) in adults with endogenous Cushing's syndrome who have type 2 diabetes mellitus or glucose intolerance and have failed surgery or cannot have surgery.

Korlym is not for people who have type 2 diabetes mellitus not caused by Cushing's syndrome.

It is not known if **Korlym** is safe and effective in children.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about Korlym?

Korlym can cause serious side effects, including:

- **Loss of a pregnancy.** Women who can become pregnant must:
 - have a negative pregnancy test before starting **Korlym** or before restarting **Korlym** if you stop taking it for more than 14 days
 - use a non-hormonal form of birth control while taking **Korlym** and for one month after stopping **Korlym**. Talk to your doctor about how to prevent pregnancy. Tell your doctor right away if you think you may be pregnant.

Please see additional Important Safety Information throughout, and [Important Facts](#) about Korlym.

There will be times
**when you visit healthcare
professionals other than the
one who prescribes Korlym for
you.** It's important for your entire
healthcare professional team to
know that you take **Korlym** and
how it may affect your care.

This guide can help you talk to your team of healthcare professionals about **Korlym** and why it was prescribed for you. It can also help you direct them to other resources so they can learn more.



IMPORTANT SAFETY INFORMATION (CONT.)

Do not take Korlym if you:

- are pregnant
- are taking medicines that contain simvastatin, lovastatin, cyclosporine, dihydroergotamine, ergotamine, fentanyl, pimozide, quinidine, sirolimus, or tacrolimus
- must take corticosteroid medicines for other serious medical problems
- have experienced unexplained vaginal bleeding, changes in the cells lining your uterus (endometrial hyperplasia), or cancer of the lining of your uterus (endometrial cancer)
- are allergic to mifepristone or any of the ingredients in **Korlym**.

Please see additional Important Safety Information throughout, and Important Facts about Korlym.

About Korlym

KEY POINTS to tell my healthcare professional team



I have endogenous hypercortisolism, also called Cushing syndrome



I have been prescribed **Korlym** to treat high blood sugar caused by endogenous hypercortisolism



Korlym works to reduce certain negative effects of excess cortisol activity



Korlym does not lower cortisol levels

My signs and symptoms of endogenous hypercortisolism may include:

- Weight gain
- High blood sugar or diabetes
- High blood pressure
- Depressed mood
- Low energy
- Rounded face
- Problems with memory and clear thinking

Your team of healthcare professionals can learn more at Korlym.com/hcp.

IMPORTANT SAFETY INFORMATION (CONT.)

What Are the Possible Side Effects of Korlym?

Serious side effects of Korlym may also include:

- **reduced effects of adrenal hormones (adrenal insufficiency).** Tell your doctor right away if you have unusual tiredness or weakness, nausea, fatigue, low blood pressure (hypotension), low blood sugar (hypoglycemia)

Please see additional Important Safety Information throughout, and [Important Facts](#) about Korlym.

Korlym
mifepristone
300 mg Tablets

Prescribing Korlym

KEY POINTS to share with my healthcare professional team



Korlym and my other prescribed medications may affect each other



Korlym is only available through Specialty Pharmacies, which will send the prescription directly to my home. **Korlym** is not available at local retail pharmacies



If any medication changes are made, please call or fax my medication list to the Specialty Pharmacist to make sure **Korlym** can safely be used with my medications

The Specialty Pharmacist team is on call 24/7 to provide support.

Contact the Specialty Pharmacy

Phone: 1-855-4KORLYM (1-855-456-7596)

Fax: 1-855-288-0601

IMPORTANT SAFETY INFORMATION (CONT.)

What Are the Possible Side Effects of Korlym?

Serious side effects of Korlym may also include (cont.):

- **low blood potassium (hypokalemia).** Your doctor should check the level of potassium in your blood before you start taking **Korlym** and while you take it. Tell your doctor if you have muscle weakness, aches or cramps, abnormal or irregular heartbeats (palpitations)

Please see additional Important Safety Information throughout, and Important Facts about Korlym.

Korlym
mifepristone
300 mg Tablets

How Korlym may affect me

KEY POINTS to tell my healthcare professional team



While I'm taking **Korlym**, my potassium, blood sugar, and blood pressure levels need to be monitored



As I continue treatment with **Korlym**, my diabetes medications may need to be adjusted

It is important for your healthcare professional team to be aware of changes and side effects that may happen during treatment with Korlym.

- **Low potassium (hypokalemia)**

Low potassium can happen at any time during treatment with **Korlym**. Symptoms of low potassium may include muscle weakness, aches, cramps, abnormal or irregular heartbeat (palpitations), constipation, and swelling of the hands or feet (edema).

- **Cortisol withdrawal**

Lower cortisol activity can lead to cortisol withdrawal effects, such as fatigue (feeling very tired), aches or weakness in your muscles or joints, nausea, and abdominal pain. These are not all of the possible side effects of **Korlym**. These symptoms may improve as you reach the best dose of **Korlym** for your body.

IMPORTANT SAFETY INFORMATION (CONT.)

What Are the Possible Side Effects of Korlym?

Serious side effects of Korlym may also include (cont.):

- **bleeding from the vagina.** **Korlym** may cause the lining of your uterus to become thick and may cause your uterus to bleed. Tell your doctor right away about any bleeding from your vagina that is not normal for you
- **disruption of menstrual cycle**

Please see additional Important Safety Information throughout, and [Important Facts](#) about Korlym.

Korlym
mifepristone
300 mg Tablets

How Korlym may affect me (cont.)

- **Blood sugar changes**

Korlym has been shown to lower blood sugar levels in some patients as early as 6 weeks.* Diabetes medications, including insulin, may need to be reduced or stopped.

- **Blood pressure changes**

Blood pressure medications may need to be increased or decreased.

- **Swelling of hands and feet (edema)**

Swelling of the hands and feet may occur during treatment with **Korlym**. If this happens along with an increase in blood pressure, it may be a sign of low potassium.

- **Vaginal bleeding**

If you have a uterus and are taking **Korlym**, you may have unusual vaginal bleeding such as spotting, missed periods, irregular periods, or other menstrual changes. If you notice signs of unusual bleeding, tell your healthcare professional team right away. It is also recommended that you notify your gynecologist that you are taking **Korlym**. Patients should not use hormonal birth control while taking **Korlym**.

- **Heart rhythm changes (QT interval prolongation)**

Korlym should not be used with certain medications or in patients with certain conditions that result in heart rhythm changes (prolonged QT interval). If treatment is necessary, the lowest effective dose should always be used.

- **Conditions treated with corticosteroids**

Korlym should not be used by patients on corticosteroid therapy for lifesaving purposes.

*In adult patients with hypercortisolism (endogenous Cushing syndrome) who had type 2 diabetes mellitus or glucose intolerance.

IMPORTANT SAFETY INFORMATION (CONT.)

What Are the Possible Side Effects of Korlym?

Serious side effects of Korlym may also include (cont.):

- problems with the electrical system of your heart (QT interval prolongation)
- worsening of symptoms of other medical problems that are treated with corticosteroids when you take corticosteroids and Korlym at the same time.

Please see additional Important Safety Information throughout, and Important Facts about Korlym.

Korlym
mifepristone
300 mg Tablets

Important information for healthcare professionals

Print out this page, fill in your information, and bring it to your team of healthcare professionals. Reprint this page anytime you need additional cards.



If you have questions about your patient's treatment, please call the prescribing physician.

Patient Name: _____

Prescriber Name: _____

Phone: _____

If you **start or change** any medications for this patient, please **call or fax the patient's medication list** to the Specialty Pharmacist for an interaction check.

Contact the Specialty Pharmacy

Phone: 1-855-4KORLYM (1-855-456-7596)

Fax: 1-855-288-0601

Healthcare professionals can report safety concerns to the 24-hour

Corcept Medical Hotline

1-855-844-3270

Korlym[®]
mifepristone
300 mg Tablets



IMPORTANT SAFETY INFORMATION (CONT.)

What Are the Possible Side Effects of Korlym? (cont.)

The most common side effects of Korlym include nausea, fatigue, headache, low potassium in your blood, pain in your arms and legs (arthralgia), vomiting, swelling of your arms and legs (peripheral edema), high blood pressure, dizziness, decreased appetite, and thickening of the lining of the uterus (endometrial hypertrophy).

These are not all the possible side effects of **Korlym**. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please see additional Important Safety Information throughout, and [Important Facts](#) about Korlym.

Korlym[®]
mifepristone
300 mg Tablets

INDICATION

Korlym is a prescription medicine used to treat high blood sugar (hyperglycemia) caused by high cortisol levels in the blood (hypercortisolism) in adults with endogenous Cushing's syndrome who have type 2 diabetes mellitus or glucose intolerance and have failed surgery or cannot have surgery.

Korlym is not for people who have type 2 diabetes mellitus not caused by Cushing's syndrome.

It is not known if **Korlym** is safe and effective in children.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about Korlym?

Korlym can cause serious side effects, including:

- **Loss of a pregnancy.** Women who can become pregnant must:
 - have a negative pregnancy test before starting **Korlym** or before restarting **Korlym** if you stop taking it for more than 14 days
 - use a non-hormonal form of birth control while taking **Korlym** and for one month after stopping **Korlym**. Talk to your doctor about how to prevent pregnancy. Tell your doctor right away if you think you may be pregnant.

Do not take Korlym if you:

- are pregnant
- are taking medicines that contain simvastatin, lovastatin, cyclosporine, dihydroergotamine, ergotamine, fentanyl, pimozide, quinidine, sirolimus, or tacrolimus
- must take corticosteroid medicines for other serious medical problems
- have experienced unexplained vaginal bleeding, changes in the cells lining your uterus (endometrial hyperplasia), or cancer of the lining of your uterus (endometrial cancer)
- are allergic to mifepristone or any of the ingredients in **Korlym**.

What Are the Possible Side Effects of Korlym?

Serious side effects of Korlym may also include:

- **reduced effects of adrenal hormones (adrenal insufficiency).** Tell your doctor right away if you have unusual tiredness or weakness, nausea, fatigue, low blood pressure (hypotension), low blood sugar (hypoglycemia)
- **low blood potassium (hypokalemia).** Your doctor should check the level of potassium in your blood before you start

taking **Korlym** and while you take it. Tell your doctor if you have muscle weakness, aches or cramps, abnormal or irregular heartbeats (palpitations)

- **bleeding from the vagina.** **Korlym** may cause the lining of your uterus to become thick and may cause your uterus to bleed. Tell your doctor right away about any bleeding from your vagina that is not normal for you
- **disruption of menstrual cycle**
- **problems with the electrical system of your heart (QT interval prolongation)**
- **worsening of symptoms of other medical problems that are treated with corticosteroids when you take corticosteroids and Korlym at the same time.**

The most common side effects of Korlym

include nausea, fatigue, headache, low potassium in your blood, pain in your arms and legs (arthralgia), vomiting, swelling of your arms and legs (peripheral edema), high blood pressure, dizziness, decreased appetite, and thickening of the lining of the uterus (endometrial hypertrophy).

These are not all the possible side effects of **Korlym**. Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Before taking Korlym, tell your doctor if you are pregnant or plan to become pregnant, have low potassium in your blood (hypokalemia), have or have had a bleeding problem or are taking medicines to thin your blood, have or have had heart problems, have had an organ transplant, have been taking medicines called corticosteroids (cortisone, dexamethasone, methylprednisolone, prednisolone, prednisone), are breastfeeding or plan to breastfeed.

You should not drink grapefruit juice while you take **Korlym**.

Tell your healthcare provider about all the medicines you take, including prescription and nonprescription medicines, vitamins, and herbal supplements. Using **Korlym** with certain other medicines may affect each other or may cause serious side effects.

Please see [Important Facts](#) about Korlym with important warning.

Korlym[®]
mifepristone
300 mg Tablets



Get 1:1 support throughout your treatment

Corcept Cares Patient Advocates are available to provide personalized support throughout treatment. They can offer educational information and resources on:

- Hypercortisolism
- How **Korlym** works
- Why monitoring is important
- What to expect during treatment

*Although **Patient Advocates** provide personalized support, they cannot offer medical advice. Talk regularly with your healthcare professional about your treatment and any side effects you may experience.*



Connect with a Patient Advocate by scanning the QR code, visiting Korlym.com, or calling 1-855-295-9969.

Please see Important Safety Information throughout, and [Important Facts](#) about Korlym, including an important warning on loss of a pregnancy.